

Datasheet for ABIN7151385
anti-ENPP1 antibody (AA 1-75)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	ENPP1
Binding Specificity:	AA 1-75
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ENPP1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Ectonucleotide pyrophosphatase/phosphodiesterase family member 1 protein (1-75AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	ENPP1
Alternative Name:	ENPP1 (ENPP1 Products)
Background:	Background: By generating PPI, plays a role in regulating pyrophosphate levels, and functions in

Target Details

bone mineralization and soft tissue calcification. PPI inhibits mineralization by binding to nascent hydroxyapatite (HA) crystals, thereby preventing further growth of these crystals. Preferentially hydrolyzes ATP, but can also hydrolyze other nucleoside 5' triphosphates such as GTP, CTP, TTP and UTP to their corresponding monophosphates with release of pyrophosphate and diadenosine polyphosphates, and also 3',5'-cAMP to AMP. May also be involved in the regulation of the availability of nucleotide sugars in the endoplasmic reticulum and Golgi, and the regulation of purinergic signaling. Appears to modulate insulin sensitivity and function.

Aliases: Alkaline phosphodiesterase 1 antibody, ARHR2 antibody, COLED antibody, E-NPP 1 antibody, Ectonucleotide pyrophosphatase/phosphodiesterase 1 antibody, Ectonucleotide pyrophosphatase/phosphodiesterase family member 1 antibody, ENPP1 antibody, ENPP1_HUMAN antibody, Ly 41 antigen antibody, M6S1 antibody, Membrane component chromosome 6 surface marker 1 antibody, NPP1 antibody, NPPase antibody, NPPS antibody, Nucleotide pyrophosphatase antibody, PC 1 antibody, PC-1 antibody, PCA1 antibody, PDNP1 antibody, Phosphodiesterase I/nucleotide pyrophosphatase 1 antibody, Plasma cell membrane glycoprotein 1 antibody, Plasma-cell membrane glycoprotein PC-1 antibody

UniProt: [P22413](#)

Pathways: [Regulation of Carbohydrate Metabolic Process](#)

Application Details

Application Notes: Recommended dilution: IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescent analysis of HepG2 cells using ABIN7151385 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)