



[Go to Product page](#)

Datasheet for ABIN7151464

## anti-EHD3 antibody (AA 269-498) (HRP)

### Overview

Quantity:	100 µg
Target:	EHD3
Binding Specificity:	AA 269-498
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EHD3 antibody is conjugated to HRP
Application:	ELISA

### Product Details

Immunogen:	Recombinant Human EH domain-containing protein 3 protein (269-498AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	EHD3
Alternative Name:	EHD3 ( <a href="#">EHD3 Products</a> )
Background:	Background: ATP- and membrane-binding protein that controls membrane reorganization/tubulation upon ATP hydrolysis (PubMed:25686250). In vitro causes tubulation

## Target Details

of endocytic membranes (PubMed:24019528). Binding to phosphatidic acid induces its membrane tubulation activity (By similarity). Plays a role in endocytic transport. Involved in early endosome to recycling endosome compartment (ERC), retrograde early endosome to Golgi, and endosome to plasma membrane (rapid recycling) protein transport. Involved in the regulation of Golgi maintenance and morphology (PubMed:16251358, PubMed:17233914, PubMed:19139087, PubMed:23781025). Involved in the recycling of internalized D1 dopamine receptor (PubMed:21791287). Plays a role in cardiac protein trafficking probably implicating ANK2 (PubMed:20489164). Involved in the ventricular membrane targeting of SLC8A1 and CACNA1C and probably the atrial membrane localization of CACNA1GG and CACNA1H implicated in the regulation of atrial myocyte excitability and cardiac conduction (By similarity). In conjunction with EHD4 may be involved in endocytic trafficking of KDR/VEGFR2 implicated in control of glomerular function (By similarity). Involved in the rapid recycling of integrin beta-3 implicated in cell adhesion maintenance (PubMed:23781025). Involved in the unidirectional retrograde dendritic transport of endocytosed BACE1 and in efficient sorting of BACE1 to axons implicating a function in neuronal APP processing (By similarity). Plays a role in the formation of the ciliary vesicle, an early step in cilium biogenesis, possibly sharing redundant functions with EHD1 (PubMed:25686250).

Aliases: EHD3 antibody, EHD2 antibody, PAST3EH domain-containing protein 3 antibody, PAST homolog 3 antibody

UniProt: [Q9NZN3](#)

## Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

## Handling

---

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.