

# Datasheet for ABIN715160 anti-Prominin 2 antibody (AA 301-400) (HRP)



$\sim$				
$O_1$	<b>/</b> el	rVI	161	Λ

Quantity:	100 μL
Target:	Prominin 2 (PROM2)
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prominin 2 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Prominin 2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Pig
Purification:	Purified by Protein A.

# Target Details

Target:	Prominin 2 (PROM2)
Alternative Name:	Prominin 2 (PROM2 Products)

## Target Details

Background:	Synonyms: PROML2, Prominin-2, PROM-2, Prominin-like protein 2, hPROML2, PROM2,
	UNQ2521/PR06014
	Background: Prominin 2 is a 112 kDa glycoporotein structurally related to Prominin 1 (CD133)
	although amino acid similarity is not more than 30 %, but their genomic organization is
	strikingly similar. Like Prominin 1, the prominin 2 exhibit similar membrane topology with 5
	trans-membrane domains and two large glycosylated extracellular domains. Similar to
	Prominin1 localization, the Prominin 2 is also associated with membrane protrusions of the
	epithelial cells from adult kidney, and all along the digestive track and other epithelial
	tissues.Prominin 2 expression is down-regulated in aggressive prostate cancer cell lines and
	transient transfection of PROML2 expression vectors has been shown to induce apoptosis in
	cultured prostate cancer cells, suggesting a tumor suppressive role for Prominin 2. Prominin 2
	expression is likely to be involved in growth suppression in the prostate, and down-regulation of
	Prominin 2 may disrupt normal prostatic homeostasis and lead to uncontrolled prostatic
	growth.
Gene ID:	150696
UniProt:	Q8N271
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish

## Handling

	peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months