antibodies -online.com





anti-NEIL1 antibody (AA 2-390)



Image



Go to Product page

Overview

Quantity:	100 μg
Target:	NEIL1
Binding Specificity:	AA 2-390
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NEIL1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Endonuclease 8-like 1 protein (2-390AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	NEIL1
Alternative Name:	NEIL1 (NEIL1 Products)
Background:	Background: Involved in base excision repair of DNA damaged by oxidation or by mutagenic
	agents. Acts as DNA glycosylase that recognizes and removes damaged bases. Has a

preference for oxidized pyrimidines, such as thymine glycol, formamidopyrimidine (Fapy) and 5-hydroxyuracil. Has marginal activity towards 8-oxoguanine. Has AP (apurinic/apyrimidinic) lyase activity and introduces nicks in the DNA strand. Cleaves the DNA backbone by beta-delta elimination to generate a single-strand break at the site of the removed base with both 3\'- and 5\'-phosphates. Has DNA glycosylase/lyase activity towards mismatched uracil and thymine, in particular in U:C and T:C mismatches. Specifically binds 5-hydroxymethylcytosine (5hmC), suggesting that it acts as a specific reader of 5hmC.

Aliases: DNA (apurinic or apyrimidinic site) Iyase Neil1 antibody, DNA glycosylase/AP Iyase Neil1 antibody, DNA-(apurinic or apyrimidinic site) Iyase Neil1 antibody, Endonuclease 8 like 1 antibody, Endonuclease 8-like 1 antibody, Endonuclease VIII antibody, Endonuclease VIII-like 1 antibody, FLJ22402 antibody, FPG1 antibody, hFPG1 antibody, NEH 1 antibody, NEH1 antibody, NEI 1 antibody, Nei endonuclease VIII like 1 (E. coli) antibody, Nei endonuclease VIII like 1 antibody, Nei homolog 1 antibody, Nei like 1 antibody, Nei like protein 1 antibody, Nei-like protein 1 antibody, NEI1 antibody, NEIL1 antibody, Neil1 antibody, NEIL1_HUMAN antibody

UniProt: Q96FI4

Pathways: DNA Damage Repair


Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human liver tissue using ABIN7151627 at dilution of 1:100