

Datasheet for ABIN7151845

anti-EPH Receptor A5 antibody (AA 246-514) (FITC)



()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Overview		
Quantity:	100 μg	
Target:	EPH Receptor A5 (EPHA5)	
Binding Specificity:	AA 246-514	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EPH Receptor A5 antibody is conjugated to FITC	
Application:	Please inquire	
Product Details		
Immunogen:	Recombinant Human Ephrin type-A receptor 5 protein (246-514AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	
Target Details		
Target:	EPH Receptor A5 (EPHA5)	
Alternative Name:	EPHA5 (EPHA5 Products)	
Background:	kground: Background: Receptor tyrosine kinase which binds promiscuously GPI-anchored ephrin- ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling in	

neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Among GPI-anchored ephrin-A ligands, EFNA5 most probably constitutes the cognate/functional ligand for EPHA5. Functions as an axon guidance molecule during development and may be involved in the development of the retinotectal, entorhino-hippocampal and hippocamposeptal pathways. Together with EFNA5 plays also a role in synaptic plasticity in adult brain through regulation of synaptogenesis. In addition to its function in the nervous system, the interaction of EPHA5 with EFNA5 mediates communication between pancreatic islet cells to regulate glucose-stimulated insulin secretion (By similarity).

Aliases: Brain-specific kinase antibody, bsk antibody, CEK 7 antibody, Eck like sequence 1 antibody, EHK 1 antibody, EHK-1 antibody, EK7 antibody, EPH homology kinase 1 antibody, EPH-like kinase 7 antibody, EPHA5 antibody, EPHA5_HUMAN antibody, Ephrin type A receptor 5 antibody, Ephrin type-A receptor 5 antibody, HEK 7 antibody, TYRO 4 antibody, Tyrosine protein kinase receptor EHK 1 antibody

UniProt: P54756

Pathways: RTK Signaling

Application Details

Restrictions: For Research Use only

Handling

Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	