

# Datasheet for ABIN7151852

# anti-EPH Receptor A7 antibody (AA 28-182) (Biotin)



#### Overview

Overview	
Quantity:	100 μg
Target:	EPH Receptor A7 (EPHA7)
Binding Specificity:	AA 28-182
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EPH Receptor A7 antibody is conjugated to Biotin
Application:	ELISA
Product Details	
Immunogen:	Recombinant Human Ephrin type-A receptor 7 protein (28-182AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified
Target Details	
Target:	EPH Receptor A7 (EPHA7)
Alternative Name:	EPHA7 (EPHA7 Products)
Background:	Background: Receptor tyrosine kinase which binds promiscuously GPI-anchored ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into

neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Among GPI-anchored ephrin-A ligands, EFNA5 is a cognate/functional ligand for EPHA7 and their interaction regulates brain development modulating cell-cell adhesion and repulsion. Has a repellent activity on axons and is for instance involved in the guidance of corticothalamic axons and in the proper topographic mapping of retinal axons to the colliculus. May also regulate brain development through a caspase(CASP3)-dependent proapoptotic activity. Forward signaling may result in activation of components of the ERK signaling pathway including MAP2K1, MAP2K2, MAPK1 AND MAPK3 which are phosphorylated upon activation of EPHA7.

Aliases: Cek 11 antibody, Developmental kinase 1 antibody, EBK antibody, EHK-3 antibody, EHK3 antibody, EK11 antibody, Embryonic brain kinase antibody, EPH homology kinase 3 antibody, EPH-like kinase 11 antibody, Epha7 antibody, EPHA7\_HUMAN antibody, Ephrin receptor Eph A7 antibody, Ephrin type A receptor 7 antibody, Ephrin type-A receptor 7 antibody, hEK11 antibody, MDK 1 antibody, Receptor protein tyrosine kinase HEK 11 antibody, Tyrosine protein kinase receptor EHK 3 antibody

UniProt: Q15375

Pathways: RTK Signaling

### Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300

Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.