antibodies -online.com







anti-EGFL7 antibody (AA 73-198)



Image



Overview

Quantity:	100 μg
Target:	EGFL7
Binding Specificity:	AA 73-198
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EGFL7 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Epidermal growth factor-like protein 7 protein (73-198AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	EGFL7
Alternative Name:	EGFL7 (EGFL7 Products)
Background:	Background: Regulates vascular tubulogenesis in vivo. Inhibits platelet-derived growth factor
	(PDGF)-BB-induced smooth muscle cell migration and promotes endothelial cell adhesion to

the extracellular matrix and angiogenesis.


Aliases: EGF like domain 7 antibody, EGF like domain containing protein 7 antibody, EGF like domain multiple 7 antibody, EGF-like protein 7 antibody, EGFL 7 antibody, EGFL7 antibody, EGFL7_HUMAN antibody, Epidermal growth factor like domain protein 7 antibody, Epidermal growth factor-like protein 7 antibody, MEGF 7 antibody, MEGF7 antibody, MGC111117 antibody, Multiple EGF like domain protein 7 antibody, Multiple EGF-like domains protein 7 antibody, Multiple epidermal growth factor like domain protein 7 antibody, Multiple epidermal growth factor-like domains protein 7 antibody, NEU1 antibody, NEU1 protein antibody, NOTCH4 like protein antibody, NOTCH4-like protein antibody, RP11 251M1.2 antibody, UNQ187/PR01449 antibody, Vascular endothelial statin antibody, VE statin antibody, VE-statin antibody, ZNEU 1 antibody, ZNEU1 antibody

UniProt:

Q9UHF1

Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7151907 diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.