antibodies .- online.com





anti-EIF3C antibody (AA 201-500) (FITC)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μg
Target:	EIF3C
Binding Specificity:	AA 201-500
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF3C antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Eukaryotic translation initiation factor 3 subunit C protein (201-500AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	EIF3C	
Alternative Name:	EIF3C (EIF3C Products)	
Background:	Background: Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates	

with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.

Aliases: Cell migration inducing protein 17 antibody, eIF 3c antibody, eIF3 p110 antibody, eIF3-p110 antibody, eIF3c antibody, EIF3C_HUMAN antibody, EIF3CL antibody, EIF3S8 antibody, EIF3S8 protein antibody, Eukaryotic translation initiation factor 3 subunit 8 (110kD) antibody, Eukaryotic translation initiation factor 3 subunit 8 110 kDa antibody, Eukaryotic translation initiation factor 3 subunit 8 antibody, Eukaryotic translation initiation factor 3 subunit C antibody

UniProt:

099613

Pathways:

Ribonucleoprotein Complex Subunit Organization

Application Details

Restrictions:

For Research Use only

Handling

Storage Comment:

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.