-online.com antibodies

Datasheet for ABIN7152214 anti-EIF5A antibody (AA 2-154) (Biotin)



\sim		•	
()	/er	110	NA
1 1	μ	$\mathbf{V} \mathbf{I} \mathbf{E}$	- 1///
\sim		V I C	~ V V

Quantity:	100 µL
Target:	EIF5A
Binding Specificity:	AA 2-154
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF5A antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Eukaryotic translation initiation factor 5A-1 protein (2-154aa)	
Isotype:	lgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	EIF5A	
Alternative Name:	EIF5A (EIF5A Products)	
Background:	Background: mRNA-binding protein involved in translation elongation. Has an important	
	function at the level of mRNA turnover, probably acting downstream of decapping. Involved in	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7152214 | 01/15/2024 | Copyright antibodies-online. All rights reserved.

	actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in
	stress response and maintenance of cell wall integrity. With syntenin SDCBP, functions as a
	regulator of p53/TP53 and p53/TP53-dependent apoptosis. Regulates also TNF-alpha-
	mediated apoptosis. Mediates effects of polyamines on neuronal process extension and
	survival. May play an important role in brain development and function, and in skeletal muscle
	stem cell differentiation. Also described as a cellular cofactor of human T-cell leukemia virus
	type I (HTLV-1) Rex protein and of human immunodeficiency virus type 1 (HIV-1) Rev protein,
	essential for mRNA export of retroviral transcripts.
	Aliases: Eukaryotic translation initiation factor 5A-1 (eIF-5A-1) (eIF-5A1) (Eukaryotic initiation
	factor 5A isoform 1) (eIF-5A) (Rev-binding factor) (eIF-4D), EIF5A
UniProt:	P63241
Pathways:	Regulation of Muscle Cell Differentiation
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.