antibodies .- online.com







anti-SLC1A6 antibody (AA 149-271) (FITC)



| () | 11/0 | K\ / | iew | 1 |
|-----|--------|------|---------------------|---|
| | \cup | ועוי | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μg |
|----------------------|--|
| Target: | SLC1A6 |
| Binding Specificity: | AA 149-271 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SLC1A6 antibody is conjugated to FITC |
| Application: | Please inquire |

Product Details

| Immunogen: | Recombinant Human Excitatory amino acid transporter 4 protein (149-271AA) | |
|-------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |

Target Details

| Target: | SLC1A6 |
|-------------------|---|
| Alternative Name: | SLC1A6 (SLC1A6 Products) |
| Background: | Background: Transports L-glutamate and also L- and D-aspartate. Seems to act as a symport by cotransporting sodium. |

Target Details

Storage Comment:

| rarget Details | |
|---------------------|--|
| | Aliases: SLC1A6 antibody, EAAT4Excitatory amino acid transporter 4 antibody, Sodium- |
| | dependent glutamate/aspartate transporter antibody, Solute carrier family 1 member 6 |
| | antibody |
| UniProt: | P48664 |
| Pathways: | Dicarboxylic Acid Transport |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 |
| | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be |
| | handled by trained staff only. |
| Storage: | -20 °C,-80 °C |

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.