Datasheet for ABIN7152295 anti-EXOSC5 antibody (AA 1-235) (FITC)

antibodies .-online.com



$\sim$				
())	$\sqrt{\rho}$	rv	ID	1/
$\cup$	v C	- I V	IC.	v v

Quantity:	100 µg
Target:	EXOSC5
Binding Specificity:	AA 1-235
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC5 antibody is conjugated to FITC
Application:	Please inquire

## Product Details

Immunogen:	Recombinant Human Exosome complex component RRP46 protein (1-235AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	EXOSC5	
Alternative Name:	EXOSC5 (EXOSC5 Products)	
Background:	Background: Non-catalytic component of the RNA exosome complex which has 3\'->5\'	
	exoribonuclease activity and participates in a multitude of cellular RNA processing and	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7152295 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Preservative:

Precaution of Use:

ProClin

	degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation
	of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing
	by-products and non-coding \'pervasive\' transcripts, such as antisense RNA species and
	promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby
	limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig
	class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by
	targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the
	RNA exosome complex is involved in general mRNA turnover and specifically degrades
	inherently unstable mRNAs containing AU-rich elements (AREs) within their 3\' untranslated
	regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems
	to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core
	complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation
	of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits
	and accessory proteins or complexes.
	Aliases: Chronic myelogenous leukemia tumor antigen 28 antibody, CML28 antibody,
	EXOS5_HUMAN antibody, EXOSC 5 antibody, Exosc5 antibody, Exosome complex component
	rrp46 antibody, exosome complex exonuclease RRP46 antibody, Exosome component 5
	antibody, exosome component Rrp46 antibody, hRrp46p antibody, p12B antibody, Ribosomal
	RNA processing protein 46 antibody, Ribosomal RNA processing protein 46, S. cerevisiae,
	homolog of antibody, Ribosomal RNA-processing protein 46 antibody, RRP41B antibody,
	RRP46 antibody, Rrp46p antibody
UniProt:	Q9NQT4
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7152295 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

Н	land	lina
	lana	miy

	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.