

Datasheet for ABIN7152356 anti-EYA1 antibody (AA 168-321) (Biotin)



Overview Quantity: 100 µg EYA1 Target: Binding Specificity: AA 168-321 Reactivity: Human Rabbit Host: Clonality: Polyclonal Conjugate: This EYA1 antibody is conjugated to Biotin Application: ELISA

Product Details

Immunogen:	Recombinant Human Eyes absent homolog 1 protein (168-321AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	EYA1
Alternative Name:	EYA1 (EYA1 Products)
Background:	Background: Functions both as protein phosphatase and as transcriptional coactivator for SIX1,
	and probably also for SIX2, SIX4 and SIX5 (By similarity). Tyrosine phosphatase that

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7152356 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Application Notes:	Optimal working dilution should be determined by the investigator.
Application Details	
Pathways:	Sensory Perception of Sound, Positive Regulation of Response to DNA Damage Stimulus
UniProt:	Q99502
	homolog 1 antibody, Eyes absent homolog1 antibody, MGC141875 antibody, OFC1 antibody
	absent 1 homolog antibody, Eyes absent homolog 1 (Drosophila) antibody, Eyes absent
	phosphatase 1 antibody, Eya1 antibody, EYA1_HUMAN antibody, Eyes absent 1 antibody, Eyes
	Aliases: BOP antibody, BOR antibody, BOS1 antibody, EYA transcriptional coactivator and
	redundant with EYA2 (By similarity).
	with SIX1, it plays an important role in hypaxial muscle development, in this it is functionally
	development of the craniofacial and trunk skeleton, kidneys and ears (By similarity). Together
	phosphorylated on Ser and Thr residues (in vitro) (By similarity). Required for normal embryonic
	during organogenesis (By similarity). Has also phosphatase activity with proteins
	Its function as histone phosphatase may contribute to its function in transcription regulation
	distinguishes between apoptotic and repair responses to genotoxic stress (PubMed:19234442).
	phosphorylation of histone H2AX plays a central role in DNA repair and acts as a mark that
	repair via the recruitment of DNA repair complexes containing MDC1. \'Tyr-142\'
	dephosphorylates \'Tyr-142\' of histone H2AX (H2AXY142ph) and promotes efficient DNA

Restrictions:	For Research Use only	

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.