antibodies .-online.com

Datasheet for ABIN7152473 anti-FBXL2 antibody (AA 1-423) (Biotin)



\sim		•	
()	er،	110	1 4 /
1 1/1			1///
$\smile v$		v i C	, v v

Quantity:	100 µg
Target:	FBXL2
Binding Specificity:	AA 1-423
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXL2 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human F-box/LRR-repeat protein 2 protein (1-423AA)	
lsotype:	lgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	FBXL2
Alternative Name:	FBXL2 (FBXL2 Products)
Background:	Background: Calcium-activated substrate recognition component of the SCF (SKP1-cullin-F-box
	protein) E3 ubiquitin-protein ligase complex, SCF(FBXL2), which mediates the ubiquitination

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7152473 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

FBL2 antibody, FBL3 antibody, FBXL 2 antibody, FBXL2 antibody, FBXL2_HUMAN antibody
FBL2/FBL3 antibody, F-box/LRR-repeat protein 2 antibody, FBL 2 antibody, FBL 3 antibody,
repeat protein 2 antibody, F-box and leucine-rich repeat protein 2 antibody, F-box protein
protein containing leucine rich repeats antibody, F box protein FBL2/FBL3 antibody, F box/LRR
Aliases: DKFZP564P0622 antibody, F box and leucine rich repeat protein 2 antibody, F box
membranes and of pulmonary surfactant (By similarity).
degradation regulates synthesis of phosphatidylcholine, which is utilized for formation of
autophagy (PubMed:23604317). PCYT1A monoubiquitination by SCF(FBXL2) and subsequent
proteasomal degradation thereby regulating phosphatidylinositol 3-kinase signaling and
(PubMed:22020328, PubMed:22323446). SCF(FBXL2) also mediates PIK3R2 ubiquitination and
calmodulin. Through CCND2 and CCND3 degradation induces cell-cycle arrest in G(0)
CCND2 and CCND3 which polyubiquitination and subsequent degradation are inhibited by
binding motifs and is thereby antagonized by calmodulin. This is the case for the cyclins
FBXL2 does not seem to target phosphodegron within its substrates but rather calmodulin-
and subsequent proteasomal degradation of target proteins. Unlike many F-box proteins,

UniProt:

Preservative:

Q9UKC9

ProClin

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.