# antibodies - online.com







# anti-FBXW5 antibody (AA 265-494)

**Images** 



$\sim$	
( )\/白	rview
OVC	

Quantity:	100 μg
Target:	FBXW5
Binding Specificity:	AA 265-494
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXW5 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

### **Product Details**

Immunogen:	Recombinant Human F-box/WD repeat-containing protein 5 protein (265-494AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	FBXW5
Alternative Name:	FBXW5 (FBXW5 Products)
Background:	Background: Substrate recognition component of both SCF (SKP1-CUL1-F-box protein) and
	DCX (DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complexes. Substrate recognition

component of the SCF(FBXW5) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of SASS6 during S phase, leading to prevent centriole reduplication. The SCF(FBXW5) complex also mediates ubiquitination and degradation of actin-regulator EPS8 during G2 phase, leading to the transient degradation of EPS8 and subsequent cell shape changes required to allow mitotic progression. Substrate-specific adapter of the DCX(FBXW5) E3 ubiquitin-protein ligase complex which mediates the polyubiquitination and subsequent degradation of TSC2. May also act as a negative regulator of MAP3K7/TAK1 signaling in the interleukin-1B (IL1B) signaling pathway.


Aliases: FBXW5 antibody, FBW5 antibody, PP3971F-box/WD repeat-containing protein 5 antibody, F-box and WD-40 domain-containing protein 5 antibody

UniProt:

Q969U6


#### **Application Details**

Application Notes:	Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded human appendix tissue using ABIN7152505 at dilution of 1:100



#### **Western Blotting**

Image 2. Western blot All lanes: FBXW5 antibody at  $2\,\mu$  g/mL + PC-3 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 64, 43 kDa Observed band size: 64 kDa



#### Immunofluorescence

**Image 3.** Immunofluorescent analysis of Hela cells using ABIN7152505 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)