# antibodies .- online.com







## anti-FANCD2 antibody (AA 1-291)

**Images** 



#### Overview

Quantity:	100 μg
Target:	FANCD2
Binding Specificity:	AA 1-291
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FANCD2 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

## **Product Details**

Immunogen:	Recombinant Human Fanconi anemia group D2 protein (1-291AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	FANCD2
Alternative Name:	FANCD2 (FANCD2 Products)
Background:	Background: Required for maintenance of chromosomal stability. Promotes accurate and
	efficient pairing of homologs during meiosis. Involved in the repair of DNA double-strand

breaks, both by homologous recombination and single-strand annealing. May participate in S phase and G2 phase checkpoint activation upon DNA damage. Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell division, particularly after replication stress. Required for the targeting, or stabilization, of BLM to non-centromeric abnormal structures induced by replicative stress. Promotes BRCA2/FANCD1 loading onto damaged chromatin. May also be involved in B-cell immunoglobulin isotype switching.

Aliases: DKFZp762A223 antibody, FA 4 antibody, FA D2 antibody, FA4 antibody, FAC D2 antibody, FACD 2 antibody, FACD 2 antibody, FACD 2 antibody, FANCD 3 antibody, FANCD 3 antibody, FANCD 3 antibody, FANCD 3 antibody, FANCONI ANEMIA COMPLEMENTATION GROUP D antibody, Fanconi anemia complementation group D2 antibody, Fanconi anemia group D2 protein antibody, FANCONI PANCYTOPENIA TYPE 4 antibody, FLJ23826 antibody, OTTHUMP00000158853 antibody, OTTHUMP00000207925 antibody, Protein FACD2 antibody, Type 4 Fanconi pancytopenia antibody

UniProt: Q9BXW9

Pathways: DNA Damage Repair

## **Application Details**

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

## Handling

Format:

Liquid

Buffer:

Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative:

ProClin

Precaution of Use:


This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

-20 °C,-80 °C


Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded human testis tissue using ABIN7152548 at dilution of 1:100



### **Immunohistochemistry**

**Image 2.** Immunohistochemistry of paraffin-embedded human brain tissue using ABIN7152548 at dilution of 1:100