

Datasheet for ABIN7152666  
**anti-FCRL5 antibody (AA 419-553)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	100 µL
Target:	FCRL5
Binding Specificity:	AA 419-553
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FCRL5 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

## Product Details

Immunogen:	Recombinant Human Fc receptor-like protein 5 protein (419-553AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	FCRL5
Alternative Name:	FCRL5 ( <a href="#">FCRL5 Products</a> )
Background:	Background: May be involved in B-cell development and differentiation in peripheral lymphoid organs and may be useful markers of B-cell stages. May have an immunoregulatory role in

## Target Details

marginal zone B-cells.

Aliases: BXMAS1 antibody, CD307 antigen antibody, CD307e antibody, Fc receptor homolog 5 antibody, Fc receptor like 5 antibody, Fc receptor-like protein 5 antibody, FcR-like protein 5 antibody, FcRH5 antibody, FcRL5 antibody, FCRL5\_HUMAN antibody, Immune receptor translocation-associated protein 2 antibody, Immunoglobulin receptor translocation associated gene 2 protein antibody, Immunoglobulin superfamily receptor translocation associated gene 2 protein antibody, IRTA2 antibody, UNQ503/PRO820 antibody

UniProt: [Q96RD9](#)

## Application Details

Application Notes: Recommended dilution: IF:1:50-1:200,

Restrictions: For Research Use only

## Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



#### Immunofluorescence

**Image 1.** Immunofluorescence staining of HeLa cells with ABIN7152666 at 1:133, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).