antibodies - online.com







anti-FRMD7 antibody (AA 464-617)

Images



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μg
Target:	FRMD7
Binding Specificity:	AA 464-617
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FRMD7 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human FERM domain-containing protein 7 protein (464-617AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	FRMD7
Alternative Name:	FRMD7 (FRMD7 Products)
Background:	Background: Plays a role in neurite development, may be through the activation of the GTPase RAC1. Plays a role in the control of eye movement and gaze stability.

Target Details

	Aliases: FRMD7FERM domain-containing protein 7 antibody
UniProt:	Q6ZUT3


Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500,	
Restrictions:	For Research Use only	

Handling


Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	

Images




Immunohistochemistry

Image 1. IHC image of ABIN7152686 diluted at 1:400 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blotting

Image 2. Western Blot Positive WB detected in: U87 whole cell lysate All lanes: FRMD7 antibody at 3.7 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 82, 80 kDa Observed band size: 82 kDa



Immunohistochemistry

Image 3. IHC image of ABIN7152686 diluted at 1:400 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.