antibodies.com

Datasheet for ABIN7152970 anti-FIZ1 antibody (AA 252-410)

2 Images



Overview

Quantity:	100 µg
Target:	FIZ1
Binding Specificity:	AA 252-410
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FIZ1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Flt3-interacting zinc finger protein 1 protein (252-410AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified


Target Details

Target:	FIZ1
Alternative Name:	FIZ1 (FIZ1 Products)
Background:	Background: May be a transcriptional repressor of NRL function in photoreceptors. Does not repress CRX-mediated transactivation (By similarity).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7152970 | 09/09/2023 | Copyright antibodies-online. All rights reserved.


Target Details	
	Aliases: FIZ1 antibody, ZNF798Flt3-interacting zinc finger protein 1 antibody, Zinc finger protein 798 antibody
UniProt:	Q96SL8
Application Details	
Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7152970 diluted at 1:300 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blotting

Image 2. Western Blot Positive WB detected in: 293 whole cell lysate All lanes: FIZ1 antibody at $4.5 \mu g/mL$ Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 52 kDa Observed band size: 52 kDa

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7152970 | 09/09/2023 | Copyright antibodies-online. All rights reserved.