

Datasheet for ABIN7153080
anti-FOXO1 antibody (AA 1-300)[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	FOXO1
Binding Specificity:	AA 1-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Forkhead box protein O1 protein (1-300AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	FOXO1
Alternative Name:	FOXO1 (FOXO1 Products)
Background:	Background: Transcription factor that is the main target of insulin signaling and regulates metabolic homeostasis in response to oxidative stress. Binds to the insulin response element (IRE) with consensus sequence 5\\\'-TT[G/A]TTTTG-3\\\' and the related Daf-16 family binding element (DBE) with consensus sequence 5\\\'-TT[G/A]TTTAC-3\\\' . Activity suppressed by

Target Details

insulin. Main regulator of redox balance and osteoblast numbers and controls bone mass. Orchestrates the endocrine function of the skeleton in regulating glucose metabolism. Acts synergistically with ATF4 to suppress osteocalcin/BGLAP activity, increasing glucose levels and triggering glucose intolerance and insulin insensitivity. Also suppresses the transcriptional activity of RUNX2, an upstream activator of osteocalcin/BGLAP. In hepatocytes, promotes gluconeogenesis by acting together with PPARGC1A and CEBPA to activate the expression of genes such as IGFBP1, G6PC and PCK1. Important regulator of cell death acting downstream of CDK1, PKB/AKT1 and SKT4/MST1. Promotes neural cell death. Mediates insulin action on adipose tissue. Regulates the expression of adipogenic genes such as PPARG during preadipocyte differentiation and, adipocyte size and adipose tissue-specific gene expression in response to excessive calorie intake. Regulates the transcriptional activity of GADD45A and repair of nitric oxide-damaged DNA in beta-cells. Required for the autophagic cell death induction in response to starvation or oxidative stress in a transcription-independent manner. Aliases: FKH 1 antibody, FKH1 antibody, FKHR antibody, Forkhead (Drosophila) homolog 1 (rhabdomyosarcoma) antibody, Forkhead box O1 antibody, Forkhead box protein O1 antibody, Forkhead box protein O1A antibody, Forkhead in rhabdomyosarcoma antibody, Forkhead, Drosophila, homolog of, in rhabdomyosarcoma antibody, FoxO transcription factor antibody, foxo1 antibody, FOXO1_HUMAN antibody, FOXO1A antibody, OTTHUMP00000018301 antibody

UniProt:	Q12778
Pathways:	PI3K-Akt Signaling , Cell Division Cycle , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Carbohydrate Homeostasis , Chromatin Binding , Regulation of Carbohydrate Metabolic Process , CXCR4-mediated Signaling Events , BCR Signaling

Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin

Handling


Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of HeLa cells using ABIN7153080 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blotting

Image 2. Western Blot Positive WB detected in: Mouse brain tissue, Mouse lung tissue All lanes: FOXO1 antibody at 2.5 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 70 kDa Observed band size: 70 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human breast cancer using ABIN7153080 at dilution of 1:100