.-online.com antibodies

# Datasheet for ABIN7153167 anti-FFAR3 antibody (AA 280-346)

4 Images



### Overview

Quantity:	100 µg
Target:	FFAR3
Binding Specificity:	AA 280-346
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FFAR3 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC)

# Product Details

Immunogen:	Recombinant Human Free fatty acid receptor 3 protein (280-346AA)
lsotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

# Target Details

Target:	FFAR3
Alternative Name:	FFAR3 (FFAR3 Products)
Background:	Background: G protein-coupled receptor that is activated by a major product of dietary fiber
	digestion, the short chain fatty acids (SCFAs), and that plays a role in the regulation of whole-

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN7153167 | 09/09/2023 | Copyright antibodies-online. All rights reserved. body energy homeostasis and in intestinal immunity. In omnivorous mammals, the short chain fatty acids acetate, propionate and butyrate are produced primarily by the gut microbiome that metabolizes dietary fibers. SCFAs serve as a source of energy but also act as signaling molecules. That G protein-coupled receptor is probably coupled to the pertussis toxin-sensitive, G(i/o)-alpha family of G proteins. Its activation results in the formation of inositol 1,4,5trisphosphate, the mobilization of intracellular calcium, the phosphorylation of the MAPK3/ERK1 and MAPK1/ERK2 kinases and the inhibition of intracellular cAMP accumulation (PubMed:12711604). Activated by SCFAs and by beta-hydroxybutyrate, a ketone body produced by the liver upon starvation, it inhibits N-type calcium channels and modulates the activity of sympathetic neurons through a signaling cascade involving the beta and gamma subunits of its coupled G protein, phospholipase C and MAP kinases. Thereby, it may regulate energy expenditure through the control of the sympathetic nervous system that controls for instance heart rate. Upon activation by SCFAs accumulating in the intestine, it may also signal to the brain via neural circuits which in turn would regulate intestinal gluconeogenesis. May also control the production of hormones involved in whole-body energy homeostasis. May for instance, regulate blood pressure through renin secretion. May also regulate secretion of the PYY peptide by enteroendocrine cells and control gut motility, intestinal transit rate, and the harvesting of energy from SCFAs produced by gut microbiota. May also indirectly regulate the production of LEP/Leptin, a hormone acting on the CNS to inhibit food intake, in response to the presence of short-chain fatty acids in the intestine. Finally, may also play a role in glucose homeostasis. Besides its role in energy homeostasis, may play a role in intestinal immunity. May mediate the activation of the inflammatory and immune response by SCFAs in the gut, regulating the rapid production of chemokines and cytokines by intestinal epithelial cells. Among SCFAs, the fatty acids containing less than 6 carbons, the most potent activators are probably propionate, butyrate and pentanoate while acetate is a poor activator (PubMed:12496283, PubMed:12711604). Aliases: FFA3R antibody, Ffar3 antibody, FFAR3\_HUMAN antibody, Free fatty acid receptor 3

antibody, G protein coupled receptor 41 antibody, G-protein coupled receptor 41 antibody, gpcr41 antibody, GPR41 antibody, gpr42 antibody

UniProt:

014843

## Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN7153167 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

# Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



#### Immunofluorescence

**Image 1.** Immunofluorescent analysis of A549 cells using ABIN7153167 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



## Immunohistochemistry

**Image 2.** Immunohistochemistry of paraffin-embedded human liver cancer using ABIN7153167 at dilution of 1:100



#### Immunofluorescence

**Image 3.** Immunofluorescent analysis of HepG2 cells using ABIN7153167 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)

Please check the product details page for more images. Overall 4 images are available for ABIN7153167.