# antibodies - online.com







## anti-GPR52 antibody (AA 225-265)



## Image



Go:	to l	Prod	luct	page
$\sim$			uot	Puge

<i>ا</i> ۱	1 /	-	rv	10	1 A
	1//	$\vdash$	1 \/	$I \leftarrow$	1//
$\sim$	v	$\sim$	1 V	-	٠٧'

Quantity:	100 μg
Target:	GPR52
Binding Specificity:	AA 225-265
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPR52 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

#### **Product Details**

Immunogen:	Recombinant Human G-protein coupled receptor 52 protein (225-265AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	GPR52
Alternative Name:	GPR52 (GPR52 Products)
Background:	Background: Orphan receptor.
	Aliases: G protein coupled receptor 52 antibody, GPR52 antibody, GPR52_HUMAN antibody,

#### **Target Details**

	Probable G protein coupled receptor 52 antibody, Probable G-protein coupled receptor 52 antibody
UniProt:	Q9Y2T5


## **Application Details**

Application Notes:	Recommended dilution: IF:1:50-1:200,
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### **Images**



#### **Immunofluorescence**

Image 1. Immunofluorescence staining of HepG2 cells with ABIN7153351 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).