



[Go to Product page](#)

Datasheet for ABIN7153705  
**anti-WDR24 antibody (AA 1-298)**

1 Image

### Overview

Quantity:	100 µL
Target:	WDR24
Binding Specificity:	AA 1-298
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDR24 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

### Product Details

Immunogen:	Recombinant Human GATOR complex protein WDR24 protein (1-298AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	WDR24
Alternative Name:	WDR24 ( <a href="#">WDR24 Products</a> )
Background:	Background: As a component of the GATOR subcomplex GATOR2, functions within the amino acid-sensing branch of the TORC1 signaling pathway (PubMed:23723238, PubMed:27166823).

## Target Details

---

Indirectly activates mTORC1 and the TORC1 signaling pathway through the inhibition of the GATOR1 subcomplex (PubMed:23723238). It is negatively regulated by the upstream amino acid sensors SESN2 and CASTOR1 (PubMed:26449471, PubMed:26586190, PubMed:27487210). In addition to its role in regulation of the TORC1 complex, promotes the acidification of lysosomes and facilitates autophagic flux (PubMed:27166823).  
Aliases: C16orf21 antibody, DKFZp434F054 antibody, JFP7 antibody, LA16c-313D11.2 antibody, WD repeat-containing protein 24 antibody, wdr24 antibody, WDR24\_HUMAN antibody

UniProt: [Q96S15](#)

## Application Details

---

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

## Handling

---

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



### Immunohistochemistry

**Image 1.** IHC image of ABIN7153705 diluted at 1:200 and staining in paraffin-embedded human ovarian tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.