

Datasheet for ABIN7153781
anti-GTF3C5 antibody (AA 305-448)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	GTF3C5
Binding Specificity:	AA 305-448
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GTF3C5 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human General transcription factor 3C polypeptide 5 protein (305-448AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	GTF3C5
Alternative Name:	GTF3C5 (GTF3C5 Products)
Background:	Background: Involved in RNA polymerase III-mediated transcription. Integral, tightly associated component of the DNA-binding TFIIIC2 subcomplex that directly binds tRNA and virus-

Target Details

associated RNA promoters.

Aliases: FLJ20857 antibody, General transcription factor 3C polypeptide 5 antibody, General transcription factor IIIC polypeptide 5 63 kDa antibody, GTF3C5 antibody, TF3C epsilon antibody, TF3C-epsilon antibody, TF3C5_HUMAN antibody, TFIIIC 63 kDa subunit antibody, TFIIIC2 63 antibody, TFIIIC63 antibody, TFIIICepsilon antibody, Transcription factor IIIC 63 kDa subunit antibody, Transcription factor IIIC epsilon subunit antibody, Transcription factor IIIC subunit epsilon antibody

UniProt: [Q9Y5Q8](#)

Application Details

Application Notes: Recommended dilution: IHC:1:20-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7153781 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.