antibodies -online.com





anti-Glucagon Receptor antibody (AA 26-136) (Biotin)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μg
Target:	Glucagon Receptor (GCGR)
Binding Specificity:	AA 26-136
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Glucagon Receptor antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Glucagon receptor protein (26-136AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	Glucagon Receptor (GCGR)
Alternative Name:	GCGR (GCGR Products)
Background:	Background: G-protein coupled receptor for glucagon that plays a central role in the regulation
	of blood glucose levels and glucose homeostasis. Regulates the rate of hepatic glucose

production by promoting glycogen hydrolysis and gluconeogenesis. Plays an important role in mediating the responses to fasting. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Promotes activation of adenylate cyclase. Besides, plays a role in signaling via a phosphatidylinositol-calcium second messenger system. Aliases: Gcgr antibody, GGR antibody, GL R antibody, GL-R antibody, GLR antibody, GLR_HUMAN antibody, Glucagon receptor antibody, Glucagon receptor precursor antibody, MGC138246 antibody

UniProt:

P47871

Pathways:

Carbohydrate Homeostasis, Regulation of Carbohydrate Metabolic Process

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.