antibodies .- online.com

Datasheet for ABIN7153872 anti-Glucagon Receptor antibody (AA 26-136) (HRP)



Overview

| Quantity: | 100 µg |
|----------------------|--|
| Target: | Glucagon Receptor (GCGR) |
| Binding Specificity: | AA 26-136 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Glucagon Receptor antibody is conjugated to HRP |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human Glucagon receptor protein (26-136AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | Glucagon Receptor (GCGR) |
|-------------------|---|
| Alternative Name: | GCGR (GCGR Products) |
| Background: | Background: G-protein coupled receptor for glucagon that plays a central role in the regulation |
| | of blood glucose levels and glucose homeostasis. Regulates the rate of hepatic glucose |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7153872 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details

| | production by promoting glycogen hydrolysis and gluconeogenesis. Plays an important role in mediating the responses to fasting. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Promotes activation of adenylate cyclase. Besides, plays a role in signaling via a phosphatidylinositol-calcium second messenger system. Aliases: Gcgr antibody, GGR antibody, GL R antibody, GL-R antibody, GLR antibody, GLR_HUMAN antibody, Glucagon receptor antibody, Glucagon receptor precursor antibody, MGC138246 antibody |
|---------------------|---|
| UniProt: | P47871 |
| Pathways: | Carbohydrate Homeostasis, Regulation of Carbohydrate Metabolic Process |
| Application Details | |
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |