

Datasheet for ABIN7154449

anti-GREM1 antibody (AA 18-184) (Biotin)



Overview

| Quantity: | 100 μg |
|----------------------|---|
| Target: | GREM1 |
| Binding Specificity: | AA 18-184 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GREM1 antibody is conjugated to Biotin |
| Application: | ELISA |

Product Details

| Immunogen: | Recombinant Human Gremlin-1 protein (18-184AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | GREM1 |
|-------------------|---|
| Alternative Name: | GREM1 (GREM1 Products) |
| Background: | Background: Cytokine that may play an important role during carcinogenesis and metanephric |
| | kidney organogenesis, as a BMP antagonist required for early limb outgrowth and patterning in |

maintaining the FGF4-SHH feedback loop. Down-regulates the BMP4 signaling in a dosedependent manner (By similarity). Antagonist of BMP2, inhibits BMP2-mediated differentiation of osteoblasts (in vitro) (PubMed:27036124). Acts as inhibitor of monocyte chemotaxis. Can inhibit the growth or viability of normal cells but not transformed cells when is overexpressed (By similarity).

Aliases: BMP antagonist 1 antibody, Cell proliferation inducing gene 2 protein antibody, Cell proliferation-inducing gene 2 protein antibody, CKTSF1B1 antibody, colorectal adenoma and carcinoma 1 antibody, Cysteine knot superfamily 1 antibody, Cysteine knot superfamily 1, BMP antagonist 1 antibody, Cysteine knot superfamily BMP antagonist 1 antibody, DAN domain family member 2 antibody, DAND2 antibody, Down regulated in Mos-transformed cells protein antibody, Down-regulated in Mos-transformed cells protein antibody, DRM antibody, grem1 antibody, GREM1_HUMAN antibody, Gremlin 1 homolog, cysteine knot superfamily antibody, Gremlin 1 like protein antibody, Gremlin 1, cysteine knot superfamily, homolog antibody, GREMLIN antibody, Gremlin-1 antibody, Gremlin1 antibody, IHG-2 antibody, IHG2 antibody, Increased in high glucose 2 antibody, Increased in high glucose protein 2 antibody, PIG2 antibody, Proliferation inducing gene 2 antibody, proliferation inducing gene 2 protein antibody

UniProt: 060565

Pathways: Regulation of Muscle Cell Differentiation, Tube Formation, Maintenance of Protein Location

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid Buffer: Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

handled by trained staff only.

-20 °C,-80 °C Storage:

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.