

Datasheet for ABIN7154450 anti-GREM1 antibody (AA 18-184) (FITC)



Overview

Quantity:	100 μg
Target:	GREM1
Binding Specificity:	AA 18-184
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GREM1 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Gremlin-1 protein (18-184AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	GREM1
Alternative Name:	GREM1 (GREM1 Products)
Background:	Background: Cytokine that may play an important role during carcinogenesis and metanephric
	kidney organogenesis, as a BMP antagonist required for early limb outgrowth and patterning in

maintaining the FGF4-SHH feedback loop. Down-regulates the BMP4 signaling in a dose-dependent manner (By similarity). Antagonist of BMP2, inhibits BMP2-mediated differentiation of osteoblasts (in vitro) (PubMed:27036124). Acts as inhibitor of monocyte chemotaxis. Can inhibit the growth or viability of normal cells but not transformed cells when is overexpressed (By similarity).

Aliases: BMP antagonist 1 antibody, Cell proliferation inducing gene 2 protein antibody, Cell proliferation-inducing gene 2 protein antibody, CKTSF1B1 antibody, colorectal adenoma and carcinoma 1 antibody, Cysteine knot superfamily 1 antibody, Cysteine knot superfamily 1, BMP antagonist 1 antibody, Cysteine knot superfamily BMP antagonist 1 antibody, DAN domain family member 2 antibody, DAND2 antibody, Down regulated in Mos-transformed cells protein antibody, Down-regulated in Mos-transformed cells protein antibody, DRM antibody, grem1 antibody, GREM1_HUMAN antibody, Gremlin 1 homolog, cysteine knot superfamily antibody, Gremlin 1 like protein antibody, Gremlin 1, cysteine knot superfamily, homolog antibody, GREMLIN antibody, Gremlin-1 antibody, Gremlin1 antibody, IHG-2 antibody, IHG2 antibody, Increased in high glucose 2 antibody, Increased in high glucose protein 2 antibody, PIG2 antibody, Proliferation inducing gene 2 protein antibody

UniProt:

060565

Pathways:

Regulation of Muscle Cell Differentiation, Tube Formation, Maintenance of Protein Location

Application Details

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.