

Datasheet for ABIN7154662

anti-VAV3 antibody (AA 562-674) (FITC)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	VAV3
Binding Specificity:	AA 562-674
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VAV3 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Human Guanine nucleotide exchange factor VAV3 protein (562-674AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	VAV3
Alternative Name:	VAV3 (VAV3 Products)
Background:	Background: Exchange factor for GTP-binding proteins RhoA, RhoG and, to a lesser extent, Rac1. Binds physically to the nucleotide-free states of those GTPases. Plays an important role

Target Details

in angiogenesis. Its recruitment by phosphorylated EPHA2 is critical for EFNA1-induced RAC1 GTPase activation and vascular endothelial cell migration and assembly (By similarity). May be important for integrin-mediated signaling, at least in some cell types. In osteoclasts, along with SYK tyrosine kinase, required for signaling through integrin alpha-v/beta-1 (ITAGV-ITGB1), a crucial event for osteoclast proper cytoskeleton organization and function. This signaling pathway involves RAC1, but not RHO, activation. Necessary for proper wound healing. In the course of wound healing, required for the phagocytotic cup formation preceding macrophage phagocytosis of apoptotic neutrophils. Responsible for integrin beta-2 (ITGB2)-mediated macrophage adhesion and, to a lesser extent, contributes to beta-3 (ITGB3)-mediated adhesion. Does not affect integrin beta-1 (ITGB1)-mediated adhesion (By similarity).

Aliases: FLJ40431 antibody, Guanine nucleotide exchange factor VAV3 antibody, Protein vav 3 antibody, Protein vav3 antibody, RGD1565941 antibody, VAV 3 antibody, Vav 3 guanine nucleotide exchange factor antibody, VAV 3 oncogene antibody, VAV 3 protein antibody, VAV-3 antibody, Vav3 antibody, VAV3 oncogene antibody, VAV3 protein antibody, VAV3_HUMAN antibody

UniProt: [Q9UKW4](#)

Pathways: [Fc-epsilon Receptor Signaling Pathway](#), [Neurotrophin Signaling Pathway](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.