antibodies -online.com





anti-GNAQ antibody (AA 1-359)

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	GNAQ
Binding Specificity:	AA 1-359
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNAQ antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Guanine nucleotide-binding protein G(q) subunit alpha protein (1-359AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	GNAQ
Alternative Name:	GNAQ (GNAQ Products)
Background:	Background: Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. Regulates B-cell selection and

survival and is required to prevent B-cell-dependent autoimmunity. Regulates chemotaxis of BM-derived neutrophils and dendritic cells (in vitro) (By similarity).

Aliases: CMC1 antibody, G alpha Q antibody, G protein alpha Q antibody, G protein subunit alpha q antibody, G-ALPHA-q antibody, GAQ antibody, GNAQ antibody, GNAQ_HUMAN antibody, guanine nucleotide binding protein (G protein), q polypeptide antibody, Guanine nucleotide binding protein alpha q antibody, guanine nucleotide binding protein G protein q polypeptide antibody, Guanine nucleotide binding protein G q subunit alpha antibody, Guanine nucleotide-binding protein alpha-q antibody, Guanine nucleotide-binding protein G(q) subunit alpha antibody, SWS antibody

UniProt:

P50148

Pathways:

JAK-STAT Signaling, Thyroid Hormone Synthesis, Myometrial Relaxation and Contraction


Application Details

Application Notes:	Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200,

Restrictions: For Research Use only


Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunofluorescence

Image 1. Immunofluorescent analysis of MCF-7 cells using ABIN7154703 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Immunohistochemistry

Image 2. Immunohistochemistry of paraffin-embedded human testis tissue using ABIN7154703 at dilution of 1:100