antibodies.com

# Datasheet for ABIN7154871 anti-HSPA1B antibody (AA 418-641)

2 Images



## Overview

Quantity:	100 µL
Target:	HSPA1B
Binding Specificity:	AA 418-641
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSPA1B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)

# Product Details

Immunogen:	Recombinant Human Heat shock 70 kDa protein 1B protein (418-641AA)
lsotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

# Target Details

Target:	HSPA1B
Alternative Name:	HSPA1B (HSPA1B Products)
Background:	Background: Molecular chaperone implicated in a wide variety of cellular processes, including
	protection of the proteome from stress, folding and transport of newly synthesized

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7154871 | 09/09/2023 | Copyright antibodies-online. All rights reserved. polypeptides, activation of proteolysis of misfolded proteins and the formation and dissociation of protein complexes. Plays a pivotal role in the protein quality control system, ensuring the correct folding of proteins, the re-folding of misfolded proteins and controlling the targeting of proteins for subsequent degradation. This is achieved through cycles of ATP binding, ATP hydrolysis and ADP release, mediated by co-chaperones. The co-chaperones have been shown to not only regulate different steps of the ATPase cycle, but they also have an individual specificity such that one co-chaperone may promote folding of a substrate while another may promote degradation. The affinity for polypeptides is regulated by its nucleotide bound state. In the ATP-bound form, it has a low affinity for substrate proteins. However, upon hydrolysis of the ATP to ADP, it undergoes a conformational change that increases its affinity for substrate proteins. It goes through repeated cycles of ATP hydrolysis and nucleotide exchange, which permits cycles of substrate binding and release. The co-chaperones are of three types: Jdomain co-chaperones such as HSP40s (stimulate ATPase hydrolysis by HSP70), the nucleotide exchange factors (NEF) such as BAG1/2/3 (facilitate conversion of HSP70 from the ADP-bound to the ATP-bound state thereby promoting substrate release), and the TPR domain chaperones such as HOPX and STUB1 (PubMed:24012426, PubMed:26865365, PubMed:24318877). Maintains protein homeostasis during cellular stress through two opposing mechanisms: protein refolding and degradation. Its acetylation/deacetylation state determines whether it functions in protein refolding or protein degradation by controlling the competitive binding of co-chaperones HOPX and STUB1. During the early stress response, the acetylated form binds to HOPX which assists in chaperone-mediated protein refolding, thereafter, it is deacetylated and binds to ubiquitin ligase STUB1 that promotes ubiquitinmediated protein degradation (PubMed:27708256). Regulates centrosome integrity during mitosis, and is required for the maintenance of a functional mitotic centrosome that supports the assembly of a bipolar mitotic spindle (PubMed:27137183). Enhances STUB1-mediated SMAD3 ubiquitination and degradation and facilitates STUB1-mediated inhibition of TGF-beta signaling (PubMed:24613385). Essential for STUB1-mediated ubiquitination and degradation of FOXP3 in regulatory T-cells (Treg) during inflammation (PubMed:23973223). Aliases: HSPA1B antibody, HSP72 antibody, Heat shock 70 kDa protein 1B antibody, Heat shock 70 kDa protein 2 antibody, HSP70-2 antibody, HSP70.2 antibody

UniProt:

#### P0DMV9

### Application Details

Application Notes:	Recommended dilution: WB:1:1000-1:5000, IF:1:50-1:200,	
Restrictions:	For Research Use only	
Order at www.antibodies-online.com   www.antiboerner-online.de   www.anticorns-enligne.fr   www.antibodies-online.cn		

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7154871 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

# Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

### Images



#### Western Blotting

**Image 1.** Western Blot Positive WB detected in: Hela whole cell lysate, A549 whole cell lysate, HepG2 whole cell lysate, Jurkat whole cell lysate, MCF-7 whole cell lysate, PC3 whole cell lysate, K562 whole cell lysate All lanes: HSPA1B antibody at 1:2000 Secondary Goat polyclonal to rabbit lgG at 1/50000 dilution Predicted band size: 71 kDa Observed band size: 71 kDa

### Immunofluorescence

**Image 2.** Immunofluorescence staining of PC3 cells with ABIN7154871 at 1:106, counter-stained with DAPI. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked in 10 % normal Goat Serum. The cells were then incubated with the antibody overnight at 4 °C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



