

Datasheet for ABIN7155047  
**anti-MST1 antibody (AA 363-465)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µg
Target:	MST1
Binding Specificity:	AA 363-465
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MST1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

## Product Details

Immunogen:	Recombinant Human Hepatocyte growth factor-like protein (363-465AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	MST1
Alternative Name:	MST1 ( <a href="#">MST1 Products</a> )
Background:	Background: extracellular exosome, extracellular region, extracellular space, receptor tyrosine kinase binding, hepatocyte growth factor receptor signaling pathway, negative regulation of

## Target Details

gluconeogenesis, regulation of cAMP-dependent protein kinase activity

Aliases: D3F15S2 antibody, DNF15S2 antibody, Hepatocyte growth factor like protein alpha chain antibody, Hepatocyte growth factor like protein antibody, Hepatocyte growth factor like protein beta chain antibody, Hepatocyte growth factor like protein homolog antibody, Hepatocyte growth factor-like protein beta chain antibody, HGFL antibody, HGFL\_HUMAN antibody, Macrophage stimulating 1 (hepatocyte growth factor like) antibody, Macrophage stimulatory protein antibody, Macrophage-stimulating protein antibody, MSP antibody, MST1 antibody, NF15S2 antibody, OTTHUMP00000208927 antibody

UniProt: [P26927](#)

## Application Details

Application Notes: Recommended dilution: IHC:1:200-1:500,

Restrictions: For Research Use only

## Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



#### Immunohistochemistry

**Image 1.** IHC image of ABIN7155047 diluted at 1:400 and staining in paraffin-embedded human liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



#### Immunohistochemistry

**Image 2.** IHC image of ABIN7155047 diluted at 1:400 and staining in paraffin-embedded human gastric cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.