antibodies - online.com







anti-HNRNPL antibody (AA 89-335)

Images



Overview

Quantity:	100 μg
Target:	HNRNPL
Binding Specificity:	AA 89-335
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPL antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Heterogeneous nuclear ribonucleoprotein L protein (89-335AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	HNRNPL
Alternative Name:	HNRNPL (HNRNPL Products)
Background:	Background: Splicing factor binding to exonic or intronic sites and acting as either an activator
	or repressor of exon inclusion. Exhibits a binding preference for CA-rich elements. Component

Target Details

of the heterogeneous nuclear ribonucleoprotein (hnRNP) complexes and associated with most nascent transcripts. Associates, together with APEX1, to the negative calcium responsive element (nCaRE) B2 of the APEX2 promoter.

Aliases: D830027H13Rik antibody, FLJ35509 antibody, Heterogeneous nuclear ribonucleoprotein L antibody, hnRNP L antibody, hnRNP-L antibody, Hnrnpl antibody, hnRPL antibody, HNRPL_HUMAN antibody, P/OKcl.14 antibody

UniProt:


P14866

Application Details

Application Notes:	Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200,	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	


Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human colon cancer using ABIN7155121 at dilution of 1:100



Western Blotting

Image 2. Western Blot Positive WB detected in: HepG2 whole cell lysate, Mouse heart tissue, Mouse brain tissue, Mouse kidney tissue All lanes: HNRNPL antibody at 3 μg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 65, 51 kDa Observed band size: 65 kDa