

Datasheet for ABIN7155213
anti-HMGB3 antibody (AA 1-103)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	HMGB3
Binding Specificity:	AA 1-103
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGB3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human High mobility group protein B3 protein (1-103AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	HMGB3
Alternative Name:	HMGB3 (HMGB3 Products)
Background:	Background: Multifunctional protein with various roles in different cellular compartments. May act in a redox sensitive manner. Associates with chromatin and binds DNA with a preference to

Target Details

non-canonical DNA structures such as single-stranded DNA. Can bent DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters (By similarity). Proposed to be involved in the innate immune response to nucleic acids by acting as a cytoplasmic promiscuous immunogenic DNA/RNA sensor (By similarity). Negatively regulates B-cell and myeloid cell differentiation. In hematopoietic stem cells may regulate the balance between self-renewal and differentiation. Involved in negative regulation of canonical Wnt signaling (By similarity).

Aliases: chromosomal protein, Nonhistone, HMG4 antibody, High mobility group (nonhistone chromosomal) protein 4 antibody, High mobility group box 3 antibody, High mobility group protein 2a antibody, High mobility group protein 4 antibody, High mobility group protein B3 antibody, High mobility group protein HMG4 antibody, HMG 4 antibody, HMG-2a antibody, HMG-4 antibody, HMG2A antibody, HMGB 3 antibody, Hmgb3 antibody, HMGB3_HUMAN antibody, MGC90319 antibody, Non histone chromosomal protein antibody, Nonhistone chromosomal protein HMG4 antibody

UniProt: [O15347](#)

Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.




Immunohistochemistry

Image 1. IHC image of ABIN7155213 diluted at 1:300 and staining in paraffin-embedded human lung cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blotting

Image 2. Western Blot Positive WB detected in: 293 whole cell lysate All lanes: HMGB3 antibody at 7 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 23 kDa Observed band size: 23 kDa



Immunohistochemistry

Image 3. IHC image of ABIN7155213 diluted at 1:300 and staining in paraffin-embedded human cervical cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.