antibodies .- online.com





anti-HIST1H2BN antibody (AA 2-126) (FITC)



Go to Product page

| \sim | |
|--------|-------|
| ()\/白 | rview |
| OVC | |

| Quantity: | 100 μg |
|----------------------|---|
| Target: | HIST1H2BN |
| Binding Specificity: | AA 2-126 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This HIST1H2BN antibody is conjugated to FITC |
| Application: | Please inquire |

Product Details

| Immunogen: | Recombinant Human Histone H2B type 1-N protein (2-126AA) |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | >95%, Protein G purified |

Target Details

| Target: | HIST1H2BN |
|-------------------|--|
| Alternative Name: | HIST1H2BN (HIST1H2BN Products) |
| Background: | Background: Core component of nucleosome. Nucleosomes wrap and compact DNA into |
| | chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a |

Target Details

template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Aliases: H2BC15 antibody, H2BFD antibody, HIST1H2BNHistone H2B type 1-N antibody, Histone H2B.d antibody, H2B/d antibody

UniProt:

Q99877

Application Details

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |