

[Go to Product page](#)

Datasheet for ABIN7155594

**anti-HLA-DQA2 antibody (AA 24-214) (Biotin)**

## Overview

Quantity:	100 µg
Target:	HLA-DQA2
Binding Specificity:	AA 24-214
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HLA-DQA2 antibody is conjugated to Biotin
Application:	ELISA

## Product Details

Immunogen:	Recombinant Human HLA class II histocompatibility antigen, DQ alpha 2 chain protein (24-214AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	HLA-DQA2
Alternative Name:	HLA-DQA2 ( <a href="#">HLA-DQA2 Products</a> )
Background:	Background: Binds peptides derived from antigens that access the endocytic route of antigen

presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accommodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases. Exogenous antigens that have been endocytosed by the APC are thus readily available for presentation via MHC II molecules, and for this reason this antigen presentation pathway is usually referred to as exogenous. As membrane proteins on their way to degradation in lysosomes as part of their normal turn-over are also contained in the endosomal/lysosomal compartments, exogenous antigens must compete with those derived from endogenous components. Autophagy is also a source of endogenous peptides, autophagosomes constitutively fuse with MHC class II loading compartments. In addition to APCs, other cells of the gastrointestinal tract, such as epithelial cells, express MHC class II molecules and CD74 and act as APCs, which is an unusual trait of the GI tract. To produce a MHC class II molecule that presents an antigen, three MHC class II molecules (heterodimers of an alpha and a beta chain) associate with a CD74 trimer in the ER to form a heterononamer. Soon after the entry of this complex into the endosomal/lysosomal system where antigen processing occurs, CD74 undergoes a sequential degradation by various proteases, including CTSS and CTSL, leaving a small fragment termed CLIP (class-II-associated invariant chain peptide). The removal of CLIP is facilitated by HLA-DM via direct binding to the alpha-beta-CLIP complex so that CLIP is released. HLA-DM stabilizes MHC class II molecules until primary high affinity antigenic peptides are bound. The MHC II molecule bound to a peptide is then transported to the cell membrane surface. In B-cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO. Primary dendritic cells (DCs) also to express HLA-DO. Lysosomal microenvironment has been implicated in the regulation of antigen loading into MHC II molecules, increased acidification produces increased proteolysis and efficient peptide loading. Aliases: DQ(6) alpha chain antibody, DQA2\_HUMAN antibody, DX alpha antibody, DX alpha chain antibody, HLA class II histocompatibility antigen antibody, HLA class II histocompatibility antigen DQ(6) alpha chain antibody, HLA class II histocompatibility antigen, DQ alpha 2 chain antibody, HLA DXA antibody, HLA-DQA1 antibody, HLA-DQA2 antibody, Major histocompatibility complex class II DQ alpha 2 antibody, MHC class II DQA2 antibody

---

UniProt: [P01906](#)

---

Pathways: [TCR Signaling, Human Leukocyte Antigen \(HLA\) in Adaptive Immune Response](#)

## Application Details

---

Application Notes: Optimal working dilution should be determined by the investigator.

---

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, -80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.