

[Go to Product page](#)

Datasheet for ABIN7156019

**anti-HIC1 antibody (AA 297-418) (HRP)**

## Overview

Quantity:	100 µg
Target:	HIC1
Binding Specificity:	AA 297-418
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIC1 antibody is conjugated to HRP
Application:	ELISA

## Product Details

Immunogen:	Recombinant Human Hypermethylated in cancer 1 protein (297-418AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	HIC1
Alternative Name:	HIC1 ( <a href="#">HIC1 Products</a> )
Background:	Background: Transcriptional repressor. Recognizes and binds to the consensus sequence \5-[CG]NG[CG]GGGCA[CA]CC-3\. May act as a tumor suppressor. May be involved in development

## Target Details

---

of head, face, limbs and ventral body wall. Involved in down-regulation of SIRT1 and thereby is involved in regulation of p53/TP53-dependent apoptotic DNA-damage responses. The specific target gene promoter association seems to be depend on corepressors, such as CTBP1 or CTBP2 and MTA1. The regulation of SIRT1 transcription in response to nutrient deprivation seems to involve CTBP1. In cooperation with MTA1 (indicative for an association with the NuRD complex) represses transcription from CCND1/cyclin-D1 and CDKN1C/p57Kip2 specifically in quiescent cells. Involved in regulation of the Wnt signaling pathway probably by association with TCF7L2 and preventing TCF7L2 and CTNNB1 association with promoters of TCF-responsive genes. Seems to repress transcription from E2F1 and ATOH1 which involves ARID1A, indicative for the participation of a distinct SWI/SNF-type chromatin-remodeling complex. Probably represses transcription from ACKR3, FGFBP1 and EFNA1.

Aliases: Hic 1 antibody, HIC ZBTB transcriptional repressor 1 antibody, Hic-1 antibody, Hic1 antibody, HIC1\_HUMAN antibody, Hypermethylated in cancer 1 antibody, Hypermethylated in cancer 1 protein antibody, ZBTB29 antibody, Zinc finger and BTB domain-containing protein 29 antibody, ZNF901 antibody

---

UniProt: [Q14526](#)

---

Pathways: [Positive Regulation of Response to DNA Damage Stimulus](#)

## Application Details

---

Application Notes: Optimal working dilution should be determined by the investigator.

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

Storage: -20 °C,-80 °C

---

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.