antibodies .- online.com





Datasheet for ABIN7156634

anti-IRF7 antibody (AA 131-289) (HRP)



Overview

Quantity:	100 μg
Target:	IRF7
Binding Specificity:	AA 131-289
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IRF7 antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Interferon regulatory factor 7 protein (131-289AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	IRF7
Alternative Name:	IRF7 (IRF7 Products)
Background:	Background: Key transcriptional regulator of type I interferon (IFN)-dependent immune
	responses and plays a critical role in the innate immune response against DNA and RNA

viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters. Can efficiently activate both the IFN-beta (IFNB) and the IFN-alpha (IFNA) genes and mediate their induction via both the virus-activated, MyD88-independent pathway and the TLR-activated, MyD88-dependent pathway. Required during both the early and late phases of the IFN gene induction but is more critical for the late than for the early phase. Exists in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, becomes phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization and nuclear localization where along with other coactivators it can activate transcription of the type I IFN and ISG genes. Can also play a role in regulating adaptive immune responses by inducing PSMB9/LMP2 expression, either directly or through induction of IRF1. Binds to the Q promoter (Qp) of EBV nuclear antigen 1 a (EBNA1) and may play a role in the regulation of EBV latency. Can activate distinct gene expression programs in macrophages and regulate the anti-tumor properties of primary macrophages.

Aliases: IMD39 antibody, Interferon regulatory factor 7 antibody, Interferon regulatory factor 7H antibody, IRF 7 antibody, IRF 7A antibody, IRF 7H antibody, IRF-7 antibody, IRF7 antibody, IRF7_HUMAN antibody, IRF7A antibody, IRF7B antibody, IRF7C antibody, IRF7H antibody

UniProt:

Q92985

Pathways:

TLR Signaling, Activation of Innate immune Response, Hepatitis C, Toll-Like Receptors Cascades, Autophagy

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.
Restrictions: For Research Use only

Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.