

Datasheet for ABIN7156707
anti-IFRD1 antibody (AA 1-200)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	IFRD1
Binding Specificity:	AA 1-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IFRD1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Interferon-related developmental regulator 1 protein (1-200AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Antigen Affinity purified & Affinity purified

Target Details

Target:	IFRD1
Alternative Name:	IFRD1 (IFRD1 Products)
Background:	Background: Could play a role in regulating gene activity in the proliferative and/or differentiative pathways induced by NGF. May be an autocrine factor that attenuates or

Target Details

amplifies the initial ligand-induced signal (By similarity).

Aliases: 12 O tetradecanoylphorbol 13 acetate induced sequence 7 antibody, IFRD1 antibody, IFRD1_HUMAN antibody, interferon related developmental regulator 1 antibody, Interferon-related developmental regulator 1 antibody, Nerve growth factor inducible protein PC4 antibody, Nerve growth factor-inducible protein PC4 antibody, PC4 antibody, Pheochromocytoma cell 4 antibody, TIS7 antibody, TPA induced sequence 7 antibody

UniProt: [O00458](#)

Application Details

Application Notes: Recommended dilution: IHC:1:500-1:5000,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7156707 diluted at 1:500 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a Goat anti-rabbit IgG labeled by HRP and visualized using 0.05 % DAB.