antibodies - online.com







anti-INF2 antibody (AA 926-1047)

Images



Overview

Quantity:	100 μg
Target:	INF2
Binding Specificity:	AA 926-1047
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This INF2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Inverted formin-2 protein (926-1047AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	INF2
Alternative Name:	INF2 (INF2 Products)
Background:	Background: Severs actin filaments and accelerates their polymerization and depolymerization.
	Aliases: C14orf151 antibody, C14orf173 antibody, CMTDIE antibody, DKFZp762A0214 antibody,

Target Details

FLJ22056 antibody, FSGS5 antibody, HBEAG binding protein 2 binding protein C antibody, HBEBP2 binding protein C antibody, HBEBP2-binding protein C antibody, INF 2 antibody, inf2 antibody, INF2_HUMAN antibody, Inverted formin 2 antibody, Inverted formin FH2 and WH2 domain containing antibody, Inverted formin-2 antibody, MGC13251 antibody, pp9484 antibody

UniProt:

Q27J81


Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7157012 diluted at 1:400 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7157012 diluted at 1:400 and staining in paraffin-embedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.