antibodies - online.com







anti-NSL1 antibody (AA 225-272)



Image



Overview

Quantity:	100 μg
Target:	NSL1
Binding Specificity:	AA 225-272
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NSL1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Recombinant Human Kinetochore-associated protein NSL1 homolog protein (225-272AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	NSL1
Alternative Name:	NSL1 (NSL1 Products)
Background:	Background: Part of the MIS12 complex which is required for normal chromosome alignment and segregation and kinetochore formation during mitosis.

Target Details

Aliases: 4833432M17Rik antibody, Al451597 antibody, C1orf48 antibody, DC31 antibody, DC8 antibody, DKFZp566O1646 antibody, Gm105 antibody, Gm984 antibody, Kinetochore associated protein NSL1 homolog antibody, Kinetochore-associated protein NSL1 homolog antibody, MIS14 antibody, NSL1 antibody, NSL1, MIND kinetochore complex component, homolog (S. cerevisiae) antibody, NSL1_HUMAN antibody

UniProt:

Q96IY1


Application Details

Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7157677 diluted at 1:300 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and



visualized using an HRP conjugated SP system.