antibodies.com

Datasheet for ABIN7158278 anti-LILRB5 antibody (AA 24-350)

Image



Overview

Quantity:	100 µL
Target:	LILRB5
Binding Specificity:	AA 24-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LILRB5 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Leukocyte immunoglobulin-like receptor subfamily B member 5 protein (24-350AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified


Target Details

Target:	LILRB5
Alternative Name:	LILRB5 (LILRB5 Products)
Background:	Background: May act as receptor for class I MHC antigens.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7158278 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
	Aliases: CD85 antigen-like family member C antibody, CD85c antibody, Leukocyte immunoglobulin-like receptor 8 antibody, Leukocyte immunoglobulin-like receptor subfamily B member 5 antibody, LILRB5 antibody, LIR-8 antibody, LIR8 antibody, LIRB5_HUMAN antibody
UniProt:	075023
Application Details	
Application Notes:	Recommended dilution: IHC:1:20-1:200,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7158278 diluted at 1:100 and staining in paraffin-embedded human ovarian cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.