antibodies -online.com





Datasheet for ABIN7158424

anti-DBT antibody (AA 140-259) (FITC)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Overview		
Quantity:	100 μg	
Target:	DBT	
Binding Specificity:	AA 140-259	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DBT antibody is conjugated to FITC	
Application:	Please inquire	
Product Details		
Product Details		
Immunogen:	Recombinant Human Lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex, mitochondrial protein (140-259AA)	
Immunogen:	dehydrogenase complex, mitochondrial protein (140-259AA)	
Immunogen: Isotype:	dehydrogenase complex, mitochondrial protein (140-259AA) IgG	
Immunogen: Isotype: Cross-Reactivity:	dehydrogenase complex, mitochondrial protein (140-259AA) IgG Human	
Immunogen: Isotype: Cross-Reactivity: Purification:	dehydrogenase complex, mitochondrial protein (140-259AA) IgG Human	
Immunogen: Isotype: Cross-Reactivity: Purification: Target Details	dehydrogenase complex, mitochondrial protein (140-259AA) IgG Human >95%, Protein G purified	

Target Details

Background:

Background: The branched-chain alpha-keto dehydrogenase complex catalyzes the overall conversion of alpha-keto acids to acyl-CoA and CO(2). It contains multiple copies of three enzymatic components: branched-chain alpha-keto acid decarboxylase (E1), lipoamide acyltransferase (E2) and lipoamide dehydrogenase (E3). Within this complex, the catalytic function of this enzyme is to accept, and to transfer to coenzyme A, acyl groups that are generated by the branched-chain alpha-keto acid decarboxylase component. Aliases: DBT antibody, BCATE2 antibody, Lipoamide acyltransferase component of branchedchain alpha-keto acid dehydrogenase complex antibody, mitochondrial antibody, EC 2.3.1.168 antibody, 52 kDa mitochondrial autoantigen of primary biliary cirrhosis antibody, Branched chain 2-oxo-acid dehydrogenase complex component E2 antibody, BCOADC-E2 antibody, Branched-chain alpha-keto acid dehydrogenase complex component E2 antibody, BCKAD-E2 antibody, BCKADE2 antibody, Dihydrolipoamide acetyltransferase component of branchedchain alpha-keto acid dehydrogenase complex antibody, Dihydrolipoamide branched chain transacylase antibody, Dihydrolipoyllysine-residue antibody, 2-methylpropanoyl)transferase antibody

UniProt:

P11182

Application Details

Restrictions:

For Research Use only

Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	