antibodies - online.com







anti-LITAF antibody (AA 1-161)





\sim				
	11/6	٦r١	/10	۱۸.

Overview		
Quantity:	100 μL	
Target:	LITAF	
Binding Specificity:	AA 1-161	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LITAF antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA	
Product Details		
Immunogen:	Recombinant Human Lipopolysaccharide-induced tumor necrosis factor-alpha factor protein (1-161AA)	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Antigen Affinity Purified	
Target Details		
Target:	LITAF	
Alternative Name:	LITAF (LITAF Products)	
Background:	Background: Probable role in regulating transcription of specific genes. May regulate through	

NFKB1 the expression of the CCL2/MCP-1 chemokine. May play a role in tumor necrosis factor alpha (TNF-alpha) gene expression.

Aliases: Lipopolysaccharide induced TNF alpha factor antibody, CMT1C antibody, FLJ38636 antibody, Lipopolysaccharide induced TNF alpha factor antibody, Lipopolysaccharide induced TNF factor antibody, Lipopolysaccharide induced tumor necrosis factor alpha factor antibody, Lipopolysaccharide-induced tumor necrosis factor-alpha factor antibody, LITAF antibody, LITAF_HUMAN antibody, LPS induced TNF alpha factor antibody, LPS-induced TNF-alpha factor antibody, MGC116698 antibody, MGC116700 antibody, MGC116701 antibody, MGC125274 antibody, MGC125275 antibody, MGC125276 antibody, p53 induced gene 7 protein antibody, PIG 7 antibody, PIG7 antibody, SIMPLE antibody, Small integral membrane protein of lysosome/late endosome antibody, TP53I7 antibody, Tumor protein p53 inducible protein 7 antibody

UniProt: Q99732

Pathways: Cellular Response to Molecule of Bacterial Origin


Application Details

Application Notes: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200,

Restrictions: For Research Use only


Handling

Format:	Liquid	
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human thymus tissue using ABIN7158438 at dilution of 1:100



Western Blotting

Image 2. Western blot All lanes: LITAFantibody at 1.91 μ g/mL + Hela whole cell lysate Secondary Goat polyclonal to rabbit lgG at 1/10000 dilution Predicted band size: 18, 24, 16 kDa Observed band size: 18 kDa



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human tonsil tissue using ABIN7158438 at dilution of 1:100