Datasheet for ABIN7158731 anti-KDM1B antibody (AA 451-544) (HRP)

antibodies .-online.com



\sim			
())ver	۲\/I	/۸/ ۵
\cup		V I '	

Quantity:	100 µg
Target:	KDM1B
Binding Specificity:	AA 451-544
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KDM1B antibody is conjugated to HRP
Application:	ELISA

Product Details

Immunogen:	Recombinant Human Lysine-specific histone demethylase 1B protein (451-544AA)	
lsotype:	lgG	
Cross-Reactivity:	Human	
Purification:	>95%, Protein G purified	

Target Details

Target:	KDM1B
Alternative Name:	KDM1B (KDM1B Products)
Background:	Background: Histone demethylase that demethylates \'Lys-4\' of histone H3, a specific tag for
	epigenetic transcriptional activation, thereby acting as a corepressor. Required for de novo DNA

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7158731 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	methylation of a subset of imprinted genes during oogenesis. Acts by oxidizing the substrate by	
	FAD to generate the corresponding imine that is subsequently hydrolyzed. Demethylates both	
	mono- and di-methylated \'Lys-4\' of histone H3. Has no effect on tri-methylated \'Lys-4\',	
	mono-, di- or tri-methylated \'Lys-9\', mono-, di- or tri-methylated \'Lys-27\', mono-, di- or tri-	
	methylated \'Lys-36\' of histone H3, or on mono-, di- or tri-methylated \'Lys-20\' of histone H4 (By similarity).	
	Aliases: amine oxidase (flavin containing) domain 1 antibody, amine oxidase, flavin containing 1	
	antibody, AOF1 antibody, bA204B7.3 antibody, C6orf193 antibody, dJ298J15.2 antibody, Flavin-	
	containing amine oxidase domain-containing protein 1 antibody, FLJ33898 antibody, FLJ34109	
	antibody, FLJ43328 antibody, KDM1B antibody, KDM1B_HUMAN antibody, LSD2 antibody,	
	lysine (K)-specific demethylase 1B antibody, Lysine-specific demethylase 2 antibody, Lysine-	
	specific histone demethylase 1B antibody, Lysine-specific histone demethylase 2 antibody	
UniProt:	Q8NB78	
Pathways:	Warburg Effect	
Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Preservative: 0.03 % Proclin 300	
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	-20 °C,-80 °C	

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.