



[Go to Product page](#)

Datasheet for ABIN7159088  
**anti-MTBP antibody (AA 567-724) (FITC)**

### Overview

Quantity:	100 µg
Target:	MTBP
Binding Specificity:	AA 567-724
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTBP antibody is conjugated to FITC
Application:	Please inquire

### Product Details

Immunogen:	Recombinant Human Mdm2-binding protein (567-724AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	MTBP
Alternative Name:	MTBP ( <a href="#">MTBP Products</a> )
Background:	Background: Inhibits cell migration in vitro and suppresses the invasive behavior of tumor cells (By similarity). May play a role in MDM2-dependent p53/TP53 homeostasis in unstressed cells.

## Target Details

---

Inhibits autoubiquitination of MDM2, thereby enhancing MDM2 stability. This promotes MDM2-mediated ubiquitination of p53/TP53 and its subsequent degradation.

Aliases: hMTBP antibody, MDM 2 binding protein antibody, MDM 2 BP antibody, MDM2 (mouse double minute 2) binding protein 104kD antibody, MDM2 (mouse double minute 2) binding protein antibody, MDM2 binding protein antibody, MDM2 BP antibody, Mdm2 transformed 3T3 cell double minute 2 p53 binding protein (mouse) binding protein 104 kDa antibody, Mdm2 transformed 3T3 cell double minute 2 p53 binding protein (mouse) binding protein antibody, Mdm2 transformed 3T3 cell double minute 2 p53 binding protein binding protein antibody, Mdm2-binding protein antibody, MDM2BP antibody, Mouse double minute 2 homolog binding protein antibody, MTBP antibody, MTBP\_HUMAN antibody

---

UniProt: [Q96DY7](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.