

Datasheet for ABIN7159140
anti-MED26 antibody (AA 372-523)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	MED26
Binding Specificity:	AA 372-523
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED26 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Mediator of RNA polymerase II transcription subunit 26 protein (372-523AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	MED26
Alternative Name:	MED26 (MED26 Products)
Background:	Background: Component of the Mediator complex, a coactivator involved in the regulated

Target Details

transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional pre-initiation complex with RNA polymerase II and the general transcription factors.

Aliases: Activator recruited cofactor 70 kDa component antibody, Activator-recruited cofactor 70 kDa component antibody, ARC 70 antibody, ARC70 antibody, Cofactor required for Sp1 transcriptional activation subunit 7 antibody, Cofactor required for Sp1 transcriptional activation, subunit 7 (70kD) antibody, Cofactor required for Sp1 transcriptional activation, subunit 7, 70 kDa antibody, CRSP 7 antibody, CRSP complex subunit 7 antibody, CRSP70 antibody, MED 26 antibody, Med26 antibody, MED26_HUMAN antibody, Mediator complex subunit 26 antibody, Mediator of RNA polymerase II transcription subunit 26 antibody, Transcriptional coactivator CRSP70 antibody

UniProt: [O95402](#)

Pathways: [Regulation of Lipid Metabolism by PPARalpha](#)

Application Details

Application Notes: Recommended dilution: IHC:1:500-1:1000,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. IHC image of ABIN7159140 diluted at 1:500 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.