antibodies .-online.com

## Datasheet for ABIN7159149 anti-MED28 antibody (AA 1-178) (Biotin)



## Overview

Quantity:	100 µg
Target:	MED28
Binding Specificity:	AA 1-178
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED28 antibody is conjugated to Biotin
Application:	ELISA

## Product Details

Immunogen:	Recombinant Human Mediator of RNA polymerase II transcription subunit 28 protein (1-178AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

## Target Details

Target:	MED28
Alternative Name:	MED28 (MED28 Products)
Background:	Background: Component of the Mediator complex, a coactivator involved in the regulated
	transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7159149 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

9	
	to convey information from gene-specific regulatory proteins to the basal RNA polymerase II
	transcription machinery. Mediator is recruited to promoters by direct interactions with
	regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation
	complex with RNA polymerase II and the general transcription factors. May be part of a
	complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton
	downstream of tyrosine kinase signaling pathways.
	Aliases: EG1 antibody, Endothelial-derived protein 1 antibody, FKSG20 antibody, Magicin
	antibody, med28 antibody, MED28_HUMAN antibody, Mediator complex subunit 28 antibody,
	Mediator of RNA polymerase II transcription subunit 28 antibody, Merlin and Grb2-interacting
	cytoskeletal protein antibody, Tumor angiogenesis marker EG-1 antibody
UniProt:	Q9H204
Pathways:	Stem Cell Maintenance, Regulation of Muscle Cell Differentiation
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should b
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Linon receipt store at -20°C or -80°C. Avoid repeated freeze

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.