antibodies -online.com





anti-Retinoid X Receptor beta antibody



Image



Overview

Quantity:	100 μL
Target:	Retinoid X Receptor beta (RXRB)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Retinoid X Receptor beta antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RXR beta	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	Retinoid X Receptor beta (RXRB)	
Alternative Name:	Rxr beta (RXRB Products)	
Background:	Synonyms: Retinoid X Receptor beta, DAUD I6, DAUDI6, H 2RIIBP, H2RIIBP, MHC class I promoter binding protein, NR2B2, RCoR 1, RCoR1, Retinoic acid receptor RXR beta, RXRB, RXRbeta.	

Background: Retinoid X receptors (RXRs) are members of the steroid/thyroid hormone receptor superfamily of nuclear receptor proteins which exert their effects by binding to specific DNA response elements, thus regulating gene expression in target cells. The RXR subfamily consists of at least three similar genes, RXR alpha, RXR beta and RXR gamma, all of which control transcription of target genes mediated by retinoids. RXR beta controls expression of many genes that respond to hormones and vitamins, including thyroid hormone, estrogen, retinoids and vitamin D. RXR beta controls a wide array of genes because of its ability to heterodimerize with other hormone receptors including the thyroid hormone receptor (THR), vitamin D receptor (VDR) and retinoic acid receptor (RAR).

Gene ID:

6257

Pathways:

Nuclear Receptor Transcription Pathway, Retinoic Acid Receptor Signaling Pathway, Steroid Hormone Mediated Signaling Pathway

Application Details

Δni	olication	Notes:
\neg	JIICALIOIT	110163.

WB: 1:100-1000, IHC-P: 1:100-500, IF(IHC-P): 1:50-200


Optimal working dilution should be determined by the investigator.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 1 % BSA, 50 % glycerol and 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded human stomach labeled with Rabbit Anti RXR beta/RXRB Polyclonal Antibody, Unconjugated (ABIN715916) at 1:200 followed by conjugation to the secondary antibody and DAB staining