antibodies -online.com





anti-Retinoid X Receptor beta antibody (Biotin)



Go to Product page

\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	Retinoid X Receptor beta (RXRB)	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Retinoid X Receptor beta antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human RXR beta	
Immunogen: Isotype:	KLH conjugated synthetic peptide derived from human RXR beta	
Isotype:	IgG	
Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat	
Isotype: Cross-Reactivity: Purification:	IgG Human, Mouse, Rat	
Isotype: Cross-Reactivity: Purification: Target Details	IgG Human, Mouse, Rat Purified by Protein A.	

RXRB, RXRbeta.

promoter binding protein, NR2B2, RCoR 1, RCoR1, Retinoic acid receptor RXR beta, RXR beta,

Background: Retinoid X receptors (RXRs) are members of the steroid/thyroid hormone receptor

superfamily of nuclear receptor proteins which exert their effects by binding to specific DNA response elements, thus regulating gene expression in target cells. The RXR subfamily consists of at least three similar genes, RXR alpha, RXR beta and RXR gamma, all of which control transcription of target genes mediated by retinoids. RXR beta controls expression of many genes that respond to hormones and vitamins, including thyroid hormone, estrogen, retinoids and vitamin D. RXR beta controls a wide array of genes because of its ability to heterodimerize with other hormone receptors including the thyroid hormone receptor (THR), vitamin D receptor (VDR) and retinoic acid receptor (RAR).

Gene ID:

6257

Pathways:

Nuclear Receptor Transcription Pathway, Retinoic Acid Receptor Signaling Pathway, Steroid Hormone Mediated Signaling Pathway

Application Details

Application Notes:

WB: 1:100-1000, IHC-P: 1:100-500

Optimal working dilution should be determined by the investigator.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months