antibodies - online.com







anti-MAGI1 antibody (AA 1234-1367)



Image



|)\/(| | | |
|------|--|--|--|
| | | | |

| Quantity: | 100 μg | |
|----------------------|--------------------------------------|--|
| Target: | MAGI1 | |
| Binding Specificity: | AA 1234-1367 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This MAGI1 antibody is un-conjugated | |
| Application: | ELISA, Immunofluorescence (IF) | |

Product Details

| Immunogen: | Recombinant Human Membrane-associated guanylate kinase, WW and PDZ domain-containing protein 1 protein (1234-1367AA) | |
|-------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | >95%, Protein G purified | |

Target Details

| Target: | MAGI1 |
|-------------------|---|
| Alternative Name: | MAGI1 (MAGI1 Products) |
| Background: | Background: May play a role as scaffolding protein at cell-cell junctions. May regulate acid- |


induced ASIC3 currents by modulating its expression at the cell surface (By similarity). Aliases: AIP 3 antibody, AIP-3 antibody, AIP3 antibody, Atrophin 1 interacting protein 3 antibody, Atrophin-1-interacting protein 3 antibody, BAI1 associated protein 1 antibody, BAI1-associated protein 1 antibody, BAIAP 1 antibody, BAIAP 1 antibody, BAP-1 antibody, BAP-1 antibody, BAP1 antibody, Gukmi1 antibody, MAGI 1 antibody, MAGI-1 antibody, MAGI1 antibody, MAGI1-HUMAN antibody, MAGI1c antibody, Membrane associated guanylate kinase inverted 1 antibody, membrane associated guanylate kinase WW and PDZ domain containing 1 antibody, Membrane-associated guanylate kinase inverted 1 antibody, TNRC 19 antibody, TNRC19 antibody, Trinucleotide repeat containing gene 19 antibody, Trinucleotide repeat-containing gene 19 protein antibody, WW and PDZ domain-containing protein 1 antibody, WW domain containing protein 3 antibody, WWP 3 antibody, WWP3 antibody

UniProt:

Q96QZ7

Application Details

| Application Notes: | Recommended dilution: IF:1:50-1:200, |
|--------------------|---|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Preservative: 0.03 % Proclin 300 Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |



Immunofluorescence

Image 1. Immunofluorescence staining of HepG2 cells with ABIN7159352 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).